

Anti-Taf7 (S. cerevisiae) antibody, rabbit serum

Storage Store at 4°C for short term. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.		• •
Storage Store at 4°C for short term. For long term storage store at '20°C. Aliquot to avoid repeated freezing and thawing. Concentration N/A	Product code	62-024
Aliquot to avoid repeated freezing and thawing. Concentration N/A	Size	100 μl
Concentration N/A	Storage	Store at 4°C for short term. For long term storage store at -20°C.
Buffer 0.1% sodium azide		Aliquot to avoid repeated freezing and thawing.
Purity Rabbit antiserum	Concentration	N/A
Immunogen Recombinant His-tagged Taf7 protein (1-214 aa) produced in E. coli	Buffer	0.1% sodium azide
Reactivity S. cerevisiae Taf7 protein Not tested with other species.	Purity	Rabbit antiserum
Reactivity S. cerevisiae Taf7 protein Not tested with other species. Special notes N/A Application 1. Western blotting (1/500) Not tested for other applications Taf3 functions as a component of the DNA-binding general transcription factor complex TFIID. Binding of TFIID to a promoter (with or without TATA element) is the initial step in pre-initiation complex (PIC) formation. TFIID plays a key role in the regulation of gene expression by RNA polymerase II through different activities such as transcription activator interaction, core promoter recognition and selectivity, TFIIA and TFIIB interaction, chromatin modification (histone acetylation by TAF1), facilitation of DNA opening and initiation of transcription. TAF7 is responsible for the recruitment of BDF1 to TATA element containing promoters.	Immunogen	Recombinant His-tagged Taf7 protein (1-214 aa) produced in <i>E. coli</i>
Not tested with other species. Special notes N/A Application 1. Western blotting (1/500) Not tested for other applications Background Taf3 functions as a component of the DNA-binding general transcription factor complex TFIID. Binding of TFIID to a promoter (with or without TATA element) is the initial step in pre-initiation complex (PIC) formation. TFIID plays a key role in the regulation of gene expression by RNA polymerase II through different activities such as transcription activator interaction, core promoter recognition and selectivity, TFIIA and TFIIB interaction, chromatin modification (histone acetylation by TAF1), facilitation of DNA opening and initiation of transcription. TAF7 is responsible for the recruitment of BDF1 to TATA element containing promoters.	Isotype	Rabbit IgG
Special notes N/A	Reactivity	S. cerevisiae Taf7 protein
Application 1. Western blotting (1/500) Not tested for other applications Taf3 functions as a component of the DNA-binding general transcription factor complex TFIID. Binding of TFIID to a promoter (with or without TATA element) is the initial step in pre-initiation complex (PIC) formation. TFIID plays a key role in the regulation of gene expression by RNA polymerase II through different activities such as transcription activator interaction, core promoter recognition and selectivity, TFIIA and TFIIB interaction, chromatin modification (histone acetylation by TAF1), facilitation of DNA opening and initiation of transcription. TAF7 is responsible for the recruitment of BDF1 to TATA element containing promoters.		Not tested with other species.
Not tested for other applications Taf3 functions as a component of the DNA-binding general transcription factor complex TFIID. Binding of TFIID to a promoter (with or without TATA element) is the initial step in pre-initiation complex (PIC) formation. TFIID plays a key role in the regulation of gene expression by RNA polymerase II through different activities such as transcription activator interaction, core promoter recognition and selectivity, TFIIA and TFIIB interaction, chromatin modification (histone acetylation by TAF1), facilitation of DNA opening and initiation of transcription. TAF7 is responsible for the recruitment of BDF1 to TATA element containing promoters.	Special notes	N/A
Taf3 functions as a component of the DNA-binding general transcription factor complex TFIID. Binding of TFIID to a promoter (with or without TATA element) is the initial step in pre-initiation complex (PIC) formation. TFIID plays a key role in the regulation of gene expression by RNA polymerase II through different activities such as transcription activator interaction, core promoter recognition and selectivity, TFIIA and TFIIB interaction, chromatin modification (histone acetylation by TAF1), facilitation of DNA opening and initiation of transcription. TAF7 is responsible for the recruitment of BDF1 to TATA element containing promoters.	Application	1. Western blotting (1/500)
complex TFIID. Binding of TFIID to a promoter (with or without TATA element) is the initial step in pre-initiation complex (PIC) formation. TFIID plays a key role in the regulation of gene expression by RNA polymerase II through different activities such as transcription activator interaction, core promoter recognition and selectivity, TFIIA and TFIIB interaction, chromatin modification (histone acetylation by TAF1), facilitation of DNA opening and initiation of transcription. TAF7 is responsible for the recruitment of BDF1 to TATA element containing promoters.		Not tested for other applications
Data Link SGD <u>S000004840</u> TAF7 / YMR227C UniProt <u>Q05021</u> (TAF7_YEAST)	Background	complex TFIID. Binding of TFIID to a promoter (with or without TATA element) is the initial step in pre-initiation complex (PIC) formation. TFIID plays a key role in the regulation of gene expression by RNA polymerase II through different activities such as transcription activator interaction, core promoter recognition and selectivity, TFIIA and TFIIB interaction, chromatin modification (histone acetylation by TAF1), facilitation of DNA opening and initiation of transcription. TAF7 is responsible for the recruitment of BDF1 to TATA element containing
Data Link SGD S000004840 TAF" / YMR227C UniProt Q05021 (TAF" / YEAST)	D. J. T. J	CCD Cooper to MARK (VARDOREC 11 : D + Ookers (MARK VE + CM)
	Data Link	SGD <u>S000004840</u> TAF7 / YMR227C UniProt <u>Q05021</u> (TAF7_YEAST)

1 / 2

PROCEDURES. NOT FOR MILITARY USE.



Data Images: 62-024 Anti-Taf7 (S. cerevisiae) antibody, rabbit serum




Fig.1 Detection of endonenous Taf7 by Western blotting using the anti-Taf7 antibody.

The antibody was used at 1/500 dilution.

Blotting was done with wet system.

Reference: This antibody was described and used in the following publication.

1. Takahata S. et al. Autonomous function of the amino-terminal inhibitorydomain of TAF1 in transcriptional regulation. Mol Cell Biol. 2004 Apr;24(8):3089-99.