

Lambda Protein Phosphatase

Product code	02-300 02-300-5
Size	20000U 5 x 20000U
Storage	-20°C ※Store at -80°C for long-term storage
Product Description	Functional full-length recombinant λ Protein Phosphatase (λ -PPase) expressed in E.coli. It is a Mn ²⁺ -dependent protein phosphatase with activity towards phosphorylated serine, threonine, tyrosine and histidine residues. It is the 221 amino-acid product of ORF221 open reading frame on bacteriophage lambda (1, 2). This product is an intact enzyme of high quality without tag.
Concentration	400 U/ μ l
Activity	one unit is defined as the amount of enzyme that hydrolyzes 1 nmole of p-nitrophenyl phosphate per minute at 30°C. Unit definition assays are performed with 50mM p-nitrophenyl phosphate in λ -PPase buffer, supplemented with 2 mM MnCl ₂ in a 50 μ l reaction.
Specific Activity	~400,000 U/mg
Component	λ -PPase: 400U/ μ l λ -PPase, 50 mM HEPES (pH 7.5), 100mM NaCl, 2 mM dithiothreitol, 0.1 mM MnCl ₂ , 0.1 mM EDTA, 50 % glycerol, 0.01 % Brij 35 (02-Lpp 50 μ l) 10 x Reaction Buffer (λ-PPase): 500mM Tris-HCl (pH 7.6), 1M NaCl, 20mM dithiothreitol, 1mM EDTA, 0.1% Brij 35 (02-Lpb 1ml) 10 x Mn²⁺: 20mM MnCl ₂ (02-Lmn 1ml)
Purity	Greater than 95% as determined by SDS-PAGE (CBB staining) that contains no detectable protease activity
Application	λ -PPase can be used to release phosphate groups from phosphorylated serine, threonine, tyrosine and histidine residues in proteins (2). It should be noted that different proteins are dephosphorylated at different rates. Optimal reaction temperature is 30°C. Inclusion of protease inhibitor cocktail and shortest incubation time is desired when assays are done with crude samples.
Data Link	UniProtKB/Swiss-Prot P03772 (PP_LAMBD)
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

Data Image : 02-300 Lambda Protein Phosphatase




Fig.1 SDS-PAGE of λ -Protein Phosphatase Phosphatase




Fig.2 Dephosphorylation of phospho -NPM1 protein by incubation with λ protein phosphatase in vitro.

HeLa cells treated with Nocodazole was lysed, and NPM1 protein was immunoprecipitated with anti-NPM1 antibody. The precipitate was suspended in 50 μ l of λ protein phosphatase reaction buffer added with 5 μ l of the protein phosphatase and incubated at 30°C for the indicated time (min). The reaction products were analyzed by western blotting.

Reference: This product has been used in the following publications.

1. Tojima T et al. Steering neuronal growth cones by shifting the imbalance between exocytosis and endocytosis. *J Neurosci*. 2014 May 21;34(21):7165-78. PMID: [24849351](https://pubmed.ncbi.nlm.nih.gov/24849351/)
2. Moriyama T et al. SUMO-modification and elimination of the active DNA demethylation enzyme TDG in cultured human cells. *Biochem Biophys Res Commun*. 2014 May 9;447(3):419-24. PMID: [24727457](https://pubmed.ncbi.nlm.nih.gov/24727457/)
3. Yang CC et al. Claspin recruits Cdc7 kinase for initiation of DNA replication in human cells. *Nature*

Commun. 2016, 7: 12135. PMID: [27401717](#)

4. Tomii S et al. Cortical Actin Alteration at the Matrix-Side Cytoplasm in Lung Adenocarcinoma Cells and Its Significance in Invasion. *Pathobiology*.2017; 84(4):171-183. PMID: [28002815](#)
5. Honda T & Inui M. PDZRN3 regulates differentiation of myoblasts into myotubes through transcriptional and posttranslational control of Id2. *J. Cell Physiol.* 2019, 234: 2963-2972. PMID: [30066954](#)

Useful References:

1. Cohen PTW & Cohen P (1989) "Discovery of a protein phosphatase activity encoded in the genome of bacteriophage λ ." *Biochem J.* **260**: 931-934 PMID:[2548489](#)
2. Zhuo S *et al* (1993) "Expression, purification, crystallization, and biochemical characterization of recombinant protein phosphatase." *J. Biol.Chem.***268**:17754-17761 PMID: [8394350](#)