

Anti-HIV-1 Gag p24 antibody, rabbit serum

| Product code | 65-004 |
|--------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|
| Size | 100 μl |
| Storage | Store 4°C for short term For long term storage store at -20°C. |
| | Aliquot to avoid repeated freezing and thawing. |
| Concentration | N/A |
| Buffer | 0.09% sodium azide |
| Purity | Rabbit antiserum |
| Immunogen | Purified full-size recombinant Gag p24 of HIV-1 subtype B (Ref 2) expressed |
| | in <i>E. coli</i> (Ref 2,3) |
| Isotype | Rabbit IgG |
| Reactivity | HIV-1 LAI strain |
| Special notes | N/A |
| | |
| Application | 1. Western blot (1/2,000~1/5,000) |
| | 2. Dot blot (1/3,000) |
| | 3. Immunoprecipitation (assay dependent) |
| | 4. ELISA (assay dependent) |
| Background | Other applications have not tested. HIV-1 Gag p24 is a capsid protein that constitutes the core of AIDS virus HIV-1 |
| Dackground | and is produced by digestion of its precursor Gag p55 by HIV-1 protease. This |
| | protein is indispensable to the reproduction of AIDS virus and constitutes an |
| | essential element for the AIDS virus particle construction (1). As this protein |
| | is detectable from the early stage of AIDS virus infection, it is used as a marker |
| | for observing the patient's condition after treatment, as it indicates the amount |
| | of virus in the blood. |
| | Using this antiserum in Western blotting, the bands of 24 kD, 55 kD and 41 kD |
| | corresponding respectively to HIV-p24 and its precursors p55 and p41 were |
| | observed in the extract of the AIDS virus infected cells (Fig. 1). |
| | |
| | |
| | |
| | |
| Data Link | GenBank: AAA44988.2 |
| Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC | |
| PROCEDURES. NOT FOR MILITARY USE. | |
| | |



Data Images: 65-004 Anti-HIV-1 Gag p24 antibody, rabbit serum




Fig.1 Detection of HIV-1 p24 and precursor proteins p55 and p41 by Western blotting using the anti p24 antibody.

Lane 1: Extract of MT4 cells

Lane2: Extract of MT4 cells infected with HIV-1(LAI strain)

The antiserum was diluted 2,500 fold

The upper bands are precursors of p55 of p24 and the processed intermediate proteins.

References

- 1. Freed EO "IV-1 gag proteins: diverse functions in the virus life cycle" *Virology* **251**:1-15 (1998) Review PMID: 9813197
- Adachi A et al "Production of acquired immunodeficiency syndrome-associated retrovirus in human nonhuman cells transfected with an infectious molecular clone" J Virol 59: 284-291(1986) PMID: 3016298
- 3. Tanaka N *et al* "A simple method for overproduction and purification of p24 Gag protein of human immunodeficiency virus type 1" *Microbiol Immunol* **36**:823-831 (1992) PMID: <u>1474933</u>
- 4. Saito A *et al* "Overproduction, purification, and diagnostic use of the recombinant HIV-1 Gag proteins, the precursor protein p55 and the processed products p17, p24, and p15" *Microbiol Immunol* **39**:473-483 (1995) PMID: 8569532

Related Products

05-005 HIV-1 Gag p24

65-023 Anti-HIV-1 Gag p24 antibody, rabbit polyclonal