

Anti-Influenza A Virus NP antibody, mouse monoclonal (C43)

Product code	65-110
Size	100 µg
Storage	-20°C
Concentration	1.0 mg/ml
Buffer	PBS- with 50% glycerol
Purity	Purified IgG fraction with protein A from hybridoma cell culture medium
Immunogen	Human Influenza A Virus (H2N2) Okada strain
Isotype	mouse IgG2ak
Reactivity	Reacts with NP of all influenza A viruses tested, including seasonal H2N2, H3N2, and avian H5N1, H5N2 and H1N1 (seasonal, pandemic and swine). No cross reactivity with influenza B viruses.
Application	<ol style="list-style-type: none"> 1. Western blotting (300~1,000 fold dilution) 2. Immunoprecipitation (100 fold dilution) 3. Immunofluorescent staining (200 fold dilution) 4. ELISA (assay dependent)
Background	<p>Influenza virus is an RNA virus, which causes influenza, and belongs to the family Orthomyxoviridae. Influenza virus is classified into three different genera, influenzavirus A, B, and C. They all have similar structures and compositions. The virions are 80-100nm in diameter and usually roughly spherical. The outer surface of the virion is made of a viral envelope containing two major glycoproteins, hemagglutinin (HA) and neuraminidase (NA). Influenzavirus A is further classified into subtypes based on the surface glycoproteins, HA and NA. Currently, there are 16 HA and 9 NA subtypes. The central core of the virion contains the viral RNA genome, which is packaged in the form of ribonucleoprotein complexes. Influenza virus nucleoprotein (NP) is a major component of the ribonucleoprotein complex and is abundantly expressed during the course of infection. It is a structural protein, which encapsidates the negative strand viral RNA and is essential for RNA transcription, replication and packaging. NP binds the PB1 and PB2 subunits of the viral RNA polymerase and the matrix protein M1, in addition to its binding to ssRNA. NP is also known to interact with variety of other macromolecules of both viral and cellular origins, and these interactions have been shown to be essential for the viral lifecycle.</p>
Data Link	UniProtKB: Influenza NP
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

Data Images: 65-110 Anti-Influenza A Virus NP antibody, mouse monoclonal (C43)




Fig.1 Immunofluorescence assay of MDCK cells derived from canine kidney cells, and A549 cells derived from human lung carcinoma cells, that were infected with H1N1 influenza virus (A/PuertoRico/8/34).

Samples were taken at 3, 9, and 24 hours post-infection. C43 antibody efficiently detected virus-infected MDCK and A549 cells as early as 3 h after infection. The cells were fixed with 4% paraformaldehyde in phosphate-buffered saline and permeabilized with 0.1% Triton X-100 in PBS. The bound antibody was visualized by a further reaction with an Alexa Fluor 488-conjugated secondary antibody.




Fig.2 Immunofluorescence assay of 293T cells expressing HA or NP of pandemic (H1N1) 2009 influenza A virus (A/Suita/1/2009).

C43 specifically recognized NP-expressing cells while a commercially available mouse anti-HA monoclonal antibody specifically recognized HA.




Fig.3. Western blotting of MDCK cells infected with H1N1 (A/PuertoRico/8/34), H5N1 (A/duck/HK/342/78), or H5N2 (A/crow/Kyoto/53/04) using C43 as a primary antibody.

Samples were collected at 3, 9, 24, and 48 hours post-infection. C43 detected NP after 3 hours post-infection and detected three different types of influenza viruses.




Fig.4. Identification of Influenza Nucleoprotein in crude extract of MDCK cells infected with Influenza A virus (H1N1) PuertoRico/8/34 using C43 monoclonal antibody.

10-20% gradient gel,

Blotting 15v, 60min (semi-dry)

Blocking overnight, 4°C

1st antibody 1/1000 dilution

2nd antibody 1/10000 dilution; rabbit polyclonal secondary antibody to mouse IgG- H & L (HRP) (ab97046; abcam). Positions of molecular size markers are shown in kDa on the left. NP size is 56 kDa according to Swiss-Prot.

References :This product has been used in the following publication

1. Mizuike R. et al. Development of Two Types of Rapid Diagnostic Test Kits To Detect the Hemagglutinin or Nucleoprotein of the Swine-Origin Pandemic Influenza A Virus H1N1. Clin Vaccine Immunol 18: 494–499 (2011) [PubMed ID: 21228147](https://pubmed.ncbi.nlm.nih.gov/21228147/) (IF)
2. Ueda M. et al. Maturation efficiency of viral glycoproteins in the ER impacts the production of

- influenza A virus. *Virus Research* 136: 91–97 (2008) [PubMed ID:18550190](#) (WB)
3. Okuno Y et al . A common neutralizing epitope conserved between the hemagglutinins of influenza A virus H1 and H2 strains. *J Virol* 67: 2552–2558 (1993) [PubMed ID:7682624](#) (IP)
 4. Sawaengsak C et al. Intranasal chitosan-DNA vaccines that protect across influenza virus subtypes. *Int J Pharm.* 2014 Oct 1;473(1-2):113-25. [PMID: 24998507](#) (WB, IF)