


## Anti-SPACA1 antibody, rabbit polyclonal, KO-Validated


<b>Product code</b>	73-062
<b>Size</b>	100 µg
<b>Storage</b>	-20°C
<b>Concentration</b>	1.0 mg/ml
<b>Buffer</b>	PBS <sup>-</sup> with 50% glycerol
<b>Purity</b>	Purified IgG fraction with protein A from rabbit antiserum.
<b>Immunogen</b>	C-terminal peptide of mouse SPACA1 protein, QSPTDIPVHEDDALSEWNE, conjugated with KLH
<b>Isotype</b>	Rabbit IgG
<b>Reactivity</b>	Mouse
<b>Special notes</b>	Validation: Knock-out mouse
<b>Application</b>	1. Western blotting (1/1,000 dilution) 2. Immunofluorescence staining (1/100~1/500 dilution)
<b>Background</b>	<p>SPACA1 is a membrane protein that localizes in the equatorial segment of spermatozoa in mammals and is reported to function in sperm-egg fusion. Disruption of Spaca1 led to the disappearance of the nuclear plate, a dense lining of the nuclear envelope facing the inner acrosomal membrane. This coincided with the failure of acrosomal expansion during spermiogenesis and resulted in the degeneration and disappearance of the acrosome in mature spermatozoa. SPACA1-deficient male mice are infertile.</p> <p>Molecular mass: Mouse; 33,343 Da with 305 amino acids</p>
<b>Data Link</b>	UniProtKB <a href="#">Q9DA48</a> mouse SPACA1, <a href="#">Gene ID 67652</a> mouse Spaca1
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

**Data Images:** 73-062 Anti-SPACA1 antibody, rabbit polyclonal, KO-Validated



**Fig.1** Western blotting of mouse testis extracts of wild-type and *Spaca1* knockout mice with anti-SPACA1 antibody.


20  $\mu$ g of Triton X-100 extracts from testes of wild type (*+/+*), *Spaca1* single (*+/-*) and double (*-/-*) knockout mice was reacted with anti-SPACA1 antibody at 1/1,000 dilution. Basigin was used as a loading control



**Fig. 2** Different forms of SPACA1 were identified in mouse testis and sperm extracts by western blot analysis with anti-SPACA1 antibody.

Anti-SPACA1 antibody was used at 1/1,000 dilution and as second antibody, goat anti-rabbit IgG conjugated with HRP (ab97051) was used at 10,000 dilution.

Several proteins associated with sperm undergo modification during maturation. Multiple bands are reflection of post-translational modifications by glycosylation and phosphorylation.



**Fig.3. Immunofluorescence staining of SPACA1 in step 7 round spermatids.**

Spaca1 protein in 7 round spermatids was reacted with anti-SPACA1 antibody and as a secondary antibody, Alexa-Fluor 488 conjugated anti-rabbit IgG antibody (green) was used. DNA was stained with Hoechst 33258.

**Reference: This antibody was described and used in the following publication.**

1. Fujihara Y. SPACA1-deficient male mice are infertile with abnormally shaped sperm heads reminiscent of globozoospermia. [Development](#). (2012) 139:3583-9. WB, IF. Free access.