

Anti-Calreticulin-3 / CALR3 / Calsperin antibody, rabbit serum

Product code	73-022
Size	100 μl
Storage	Store 4°C for short term For long term storage store at -20°C.
Biorage	Aliquot to avoid repeated freezing and thawing.
Concentration	N/A
Buffer	0.1% sodium azide
Purity	Rabbit antiserum
Immunogen	Synthetic peptide corresponding to C-terminal region of mouse CALR3,
	CMGKFHRHNHLSRFHRQGEL.
Isotype	Rabbit IgG
Reactivity	Mouse.
	Not tested with other species.
Special notes	N/A
Application	1. Western blotting (1/1,000 dilution))
	2. Immunoprecipitation (1/100 dilution)
	3. Immunohistochemistry (1/100 dilution)
	Other applications have not been tested.
Background	CALR3 capacity for calcium-binding may be absent or much lower than that of
	CALR. During spermatogenesis, may act as a lectin-independent chaperone for
	specific client proteins such as ADAM3. Required for sperm fertility.
	Molecular mass: 44,232 with 380 amino acids.
Data Link	uniprot/Q9D9Q6 mouse, Gene ID 73316 mouse
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC	

PROCEDURES. NOT FOR MILITARY USE.



Data Images: 73-022 Anti-Calreticulin-3 / CALR3 / Calsperin antibody, rabbit serum




Fig. 1. Testeis specific expression of Calreticulon-3 as examined in various tissues by western blotting with anti-CALR3 antibody.

The various tissues were excised and homogenized in lysis buffer containing 1% TritonX100 and then placed on ice for 1 h. These extracts were centrifuged, and the supernatants were collected and analyzed by western blotting with anti-CALR3 antibody at 1/1,000 dilution.

1.Brain. 2. Lung. 3. Heart. 4. Thymus. 5. Liver. 6. Spleen. 7. Kidney. 8. Muscle. 9. Ovary. 10. Testis. 11. Sperm




Fig.2. Identification of Calreticulon-3 protein by western blotting with anti-CALR3 antibody.

Embryonic fibroblast cells prepared from Calr3-/-mouse were transfected with a plasmid expressing Calr3. The cell lysate was analyzed by western blotting with anti-CALR3 antibody at 1/1,000 dilution.

- 1. Mock-infected cell lysate as a negative control.
- 2. Cell lysate transfected with a plasmid expressing Calr3.






Fig.3. Analysis of Calrecticulon-3 protein in the extracts of mouse testis and sperm by western blotting with anti-Calrecticulon-3 antibody.

Proteins in the extracts (10 µg protein) were separated on SDS-PAGE (10-20% gradient gel), electro-blotted to PVDF membrane and reacted with anti-Calrecticulon-3 antibody at 1/1,000 dilution. As the second antibody, anti-rabbit IgG antibody conjugated with HRP (Abcam:ab97051) was used at 1/10,000 dilution. The numbers on the left are positions of protein size markers in kDa.




Fig.4. Immunoprecipitation of Calreticulon-3 protein with anti-CALR3 antibody.

Lysates of wild-type mouse testis were immunoprecipitated with anti-CALR3 antibody and the precipitates were analyzed by western blotting with the same antibody.

- 1. Input testis lysate
- 2. Precipitated with preimmune serum
- 3. Precipitated with anti-CALR3 antibody






Fig.5 Immunofluorescence staining of a testicular section with anti-CALR3 antibody.

Sequential sections were stained with hematoxylin and eosin (upper panel). CALR3 was detected in elongating spermatids (lower panel). Testis was collected from adult mouse and fixed in 4% paraformaldehyde/PBS overnight at 4 °C, cryopreserved in graded 10-30% sucrose, and embedded. Frozen sections (8 µm) were mounted on aminopropyltriethoxysilane-coated glass slides. Primary antibody was used at 1/100 and as secondary antibody, Alexa Fluor 488 conjugated goat anti-rabbit IgG was used. Scale bar is 200 µm.

Reference: This antibody was described and used in the following publication.

1. Ikawa M. et al (2011) Calsperin is a testis-specific chaperone required for sperm fertility. J Biol <u>Chem.</u>18:5639-46. <u>pubmed/21131354</u> **WB, IP, IHC.** Free article.