


## Anti-EDD/UBR5 antibody, rabbit polyclonal

<b>Product code</b>	70-501
<b>Size</b>	100 µg
<b>Storage</b>	-20°C
<b>Concentration</b>	1.0 mg/ml
<b>Buffer</b>	PBS <sup>-</sup> with 50% glycerol
<b>Purity</b>	Affinity-purified with immunogen.
<b>Immunogen</b>	Synthetic peptide of human EDD protein corresponding to 394-408 amino acids, C-KWSESEPYRNAQNPS, conjugated with KLH
<b>Isotype</b>	Rabbit IgG
<b>Reactivity</b>	Human (HeLa, HEK293T, MCF7) and mouse (NIH3T3). Note that expression level of EDD varies greatly, depending on tissues and cell types (see Fig.1).
<b>Special notes</b>	N/A
<b>Application</b>	1. Western blotting (1/1,000) 2. Immunofluorescent staining 1/1,000)
<b>Background</b>	EDD / UBR5 (2,799 aa, 309 kDa) is E3 ubiquitin-protein ligase which is a component of the N-end rule pathway. Recognizes and binds to proteins bearing specific N-terminal residues that are destabilizing according to the N-end rule, leading to their ubiquitination and subsequent degradation. By similarity. Involved in maturation and/or transcriptional regulation of mRNA by activating CDK9 by polyubiquitination. May play a role in control of cell cycle progression. May have tumor suppressor function. Regulates DNA topoisomerase II binding protein (TopBP1) in the DNA damage response. Plays an essential role in extraembryonic development. Ubiquitinates acetylated PCK1. Also acts as a regulator of DNA damage response by acting as a suppressor of RNF168, an E3 ubiquitin-protein ligase that promotes accumulation of 'Lys-63'-linked histone H2A and H2AX at DNA damage sites, thereby acting as a guard against excessive spreading of ubiquitinated chromatin at damaged chromosomes.
<b>Data Link</b>	<a href="https://www.uniprot.org/entry/O95071">uniprot/O95071</a> UBR5 human
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

**Data Images:** 70-501 Anti-EDD/UBR5 antibody, rabbit polyclonal



**Fig.1 Identification of EDD in whole cell extracts of human cells by western blotting using anti-EDD antibody.**


Lane 1; HeLa cells transfected with control siRNA

Lane 2; HeLa cells transfected with EDD-specific siRNA

Lane 3; HEK293T cells transfected with pFlag-CMV-5b empty vector

Lane 4; HEK293T cells transfected with pCMV-Tag2b-EDD expression vector

Predicted molecular mass of EDD is 309 kDa and the corresponding band in Lane 1 is much reduced in Lane 2 by introduction of the EDD-specific siRNA. Note that expression level of EDD in HEK293T cells is much higher than in HeLa cells. Mol Mass of EDD is 309 kDa



**Fig.2. Immunofluorescence staining of EDD protein in MCF7 cells.**

MCF7 cells were fixed in 4% paraformaldehyde overnight and permeabilized

in 0.25% TritonX 100 in PBS for 10 min. Anti-EDD antibody was used at 1/1,000 dilution.

As second antibody, goat anti-rabbit IgG conjugated with Alex488 was used at 1/1,000 dilution. DNA was stained with DAPI. Note that not all nuclei were stained, indicating cell cycle dependency of EDD protein existence.

**Reference:** This antibody has been used in the following publications.

1. Watanabe S et al. GRWD1 regulates ribosomal protein L23 levels via the ubiquitin-proteasome system. [J Cell Sci](#). 2018 Aug 3;131(15). PMID:[29991511](#) **WB (human)**