

Anti- Legionella pneumophila antibody, rabbit polyclonal

Product code	64-100
Size	200 µl
Storage	Store 4°C for short term For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Concentration	N/A
Buffer	0.09% sodium azide
Purity	Rabbit antiserum
Immunogen	Formaldehyde treated whole cells of Legionella pneumophila strain Philadelphia 1 (ATCC #33152). Immunized 7 times at two weeks intervals.
Isotype	Rabbit IgG
Reactivity	Legionella pneumophila strains. Since the antiserum has not been adsorbed, it may cross-reacts with related bacteria.
Special notes	N/A
Application	<ol style="list-style-type: none"> 1. Immunofluorescent and Immunochemical staining (1/10,000 ~1/30,000 dilution) 2. Immunohistochemistry (1/3,000~1/10,000) 3. ELISA (1/10,000~1/30,000 dilution) 4. Agglutination (1/2,000~1/5,000)
Background	<p>Legionnaires disease (LD) was recognized in 1976 after an outbreak of pneumonia at an American Legion convention in Philadelphia. Soon after, the etiologic agent was identified as a fastidious gram-negative bacillus and named <i>Legionella pneumophila</i>. Although several other species of the genus <i>Legionella</i> were subsequently identified, <i>L pneumophila</i> is the most frequent cause of human legionellosis and a relatively common cause of community-acquired and nosocomial pneumonia in adults. In children, <i>L pneumophila</i> is also an important, although relatively uncommon, cause of pneumonia.</p>
Data Link	ATCC #33152
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

Data Images: 64-100 Anti- Legionella pneumophila antibody, rabbit polyclonal




Fig.1 Immunofluorescent staining of Legionella pneumophila in the infected HEK293 cells using anti-Legionella pneumophila antibody.

HEK293 cells were infected with Legionella pneumophila strain Philadelphia1 fixed with 4% formaldehyde and reacted with the anti-Legionella pneumophila antibody at 1/10,000 dilution. As the second antibody, goat Rodamine Red X conjugated anti-rabbit IgG antibody was used at 1/10,000 dilution. DNA was stained with Hoechst 33342 (center) and the images were merged with that of differential interference contrast microscope (right).




Fig.2. Titration of antibody reactivity of anti-Legionella pneumophila antiserum by direct ELISA
 Plate was coated with 100 μ l of 10^9 cells/ml per well and 100 μ l of the antiserum at the indicated dilution was added to each well and incubated. After washing, goat anti-rabbit-IgG conjugated with HRP was added as 2nd antibody. Color was developed with TMB as substrate. “pre” is preimmune serum and “Immunized” is immunized serum.