

Anti-Helicobacter pylori flagellin antibody, mouse monoclonal (hp-01)

64-108 100 μg

Shipping and Storage: Shipped at 4° C or -20° C and store at -20° C

Immunogen: Crude extract of *Helicobacter pylori*

Specific Reactivity: Reacts with Helicobacter pylori flagellin (FlaA and FlaB) protein

Isotype: mouse $IgG1 \kappa$

Product: 1 mg/ml in PBS- with 50% glycerol, filter sterilized.

Purity: purified with Protein A

Applications:

1. Western blotting (1/500~1/1,000)

2. ELISA (assay dependent)

Other applications have not been tested.


Background: Helicobacter pylori (H. pylori) is a gram-negative, spiral-shaped microaerophilic bacterium which is linked to the development of duodenal ulcers and stomach cancer. H. pylori strains produce important virulence factor. Motility is conferred to the bacteria by several sheathed flagella and is regarded as one of their principal virulence factors. Characterization of the proteins associated with the flagella components indicate the hook protein (FlgE) is 78 kDa in size while the filament consists of two flagellin proteins, FlaA and FlaB. The H. pylori flaA and flaB genes both coded for proteins with 510 and 514 amino acids and molecular masses of 53 kDa and 54 kDa, respectively. Both flagellins share considerable amino acid homology (58% identity).

Data Link: UniProt KB: <u>P0A0S1</u> (FLAA/HELPT), Flagellin A, Q07911 (FLAB/HELPT) ,Flagellin B

Fig.1. Detection of polypeptide of *H. pylori* by Western blotting with MAb (hp-01).

- 1: Crude extract of *H. pylori* (isolated strains)
- 2: Crude extract of *H.pylori* (JCM12093)
- 3: Crude extract of Campylobacter jejuni
- 4: Crude extract of Campylobacter coli
- 5: Crude extract of *Escherichia coli (E.coli)*The hp-01 antibody was used at 1/500 dilution. The HRPO-conjugate goat anti-mouse IgG was used at 1/4,000 as the second antibody and visualized by DAB (3,3'-

Diaminobenzidine). The hp-01 reacted with 53



kDa protein in the extract of *H. pylori*, but did not react with the extracts of *Campylobacter spp.* and *E. coli*. A protein band 53 kDa in size corresponds to the expected size of the flagellin.



Fig.2. Detection of FlaA and FlaB protein of *H. pylori* by Western blotting with hp-01

- 1: Crude extract of *H.pylori* (JCM12093)
- 2: Recombinant FlaA protein expressed in *E. coli*
- 3: Recombinant FlaB protein expressed in *E. coli*
- 4: Negative control (BL21(DE3))

The hp-01 antibody was used at 1/500 dilution. The HRPO-conjugate goat anti-mouse IgG was used at 1/4,000 as the second antibody and visualized by Chemi-Luminescence. The hp-01 reacted with

recombinant FlaA and FlaB proteins expressed in E. coli.





Fig.3. Titration of flagellin in the extract of *H. pylori* cells by indirect ELISA using hp-01.

The wells of plate were coated indicated amounts of the extract of H. pylori cells per well. After blocking with 5% skim milk, 100 μ l of hp-01 at 1/500 dilution was added to the each well. The HRPO-conjugated goat anti-mouse IgG (100 μ l, x2000 dilution) was added. Color was developed with OPD (orthophenylenediamine) as substrate. Optical density (OD) measured at 490nm.



| | ELISA | WB(kDa) |
|--------------------------------------|-------|---------|
| Helicobacter pylori (JCM12093) | + | 53 |
| Helicobacter pylori (England strain) | + | 53 |
| Helicobacter pylori (MR31 strain) | + | 53 |
| Helicobacter pylori (RD26 strain) | + | 53 |
| Campylobacter jejuni | _ | _ |
| Campylobacter coli | _ | _ |
| Campylobacter fetus | _ | _ |
| Escherichia coli | _ | _ |

Table 1. Reactivities of MAb (hp-01)

The hp-01 reacted with a standard strain of *H. pylori* (JCM12093) and isolated strains, such as England strain, USA strains (MR31 and RD26 strains). However hp-01 did not react with other food poisoning bacteria such as *Campylobacter jejuni*, *Campylobacter coli* and enterotoxigenic *Escherichia coli* (ETEC).

Reference: There has been no publication using this antibody.

Please let us know when your research using this antibody is published. We will offer one vial of our antibody as compliment.

Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.