

Anti-Dlx5 antibody, guinea pig serum

Product code	74-117
Size	50 µl
Storage	Store 4°C for short term For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Concentration	N/A
Buffer	0.09% sodium azide
Purity	Guinea pig antiserum
Immunogen	Purified recombinant MBP-mouse Dlx5 (amino acids 1-135) fusion protein.
Isotype	Guinea pig IgG
Reactivity	Reacts with mouse. May also react with rat and human from sequence homology.
Special notes	N/A
Application	1. Western blot (1/1,000) 2. Immunohistochemistry –Frozen Section (1/3,000)
Background	Transcriptional factor involved in bone development. Acts as an immediate early BMP-responsive transcriptional activator essential for osteoblast differentiation. Stimulates ALPL promoter activity in a RUNX2-independent manner during osteoblast differentiation. Stimulates SP7 promoter activity during osteoblast differentiation. Promotes cell proliferation by up-regulating MYC promoter activity. Involved as a positive regulator of both chondrogenesis and chondrocyte hypertrophy in the endochondral skeleton. Binds to the homeodomain-response element of the ALPL and SP7 promoter. Binds to the MYC promoter. Requires the 5'-TAATTA-3' consensus sequence for DNA-binding. Length: 289 amino acids. Mass (Da): 31,396 Subcellular localization: Nucleus.
Data Link	UniProtKB P70396 (DLX5_MOUSE)
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

Data Images: 74-117 Anti-Dlx5 antibody, guinea pig serum




Fig 1. Immunohistochemical staining of Dlx5 in mouse embryonic forebrain.

A. 13.5 forebrain cryo-section was immunostained for Dlx5.

B. Enlarged image of A of the septum. Double staining of Dlx5 with Necdin. Arrowheads indicate double-stained cells.

The anti-Dlx5 antibody was used at 1/3,000 dilution and anti-Necdin antibody (BioAcademia 74-100) was at 1/1,000.




Fig 2. Immunohistochemical staining of Dlx5 in mouse embryonic forebrain: Comparison with other markers such as Oligo2 and Lhx6, expressed in the similar region.

Sample is E13.5 Sections. The Dlx5 antiserum was used at 1/3,000 (green), anti-Oligo2 rabbit antibody (Red, Millipore) at 1/5,000, and anti-Lhx6 mouse antibody (Blue, Santa Cruz) at 1/200.

Reference: This antibody has been used in the following publications.

1. Kuwajima T et al. Necdin promotes GABAergic neuron differentiation in cooperation with Dlx homeodomain proteins. *J Neurosci.* 2006 May 17;26(20):5383-92. PMID: [16707790](https://pubmed.ncbi.nlm.nih.gov/16707790/) **WB, IHC-F (mouse)**
2. Bluske KK et al. β -Catenin signaling specifies progenitor cell identity in parallel with Shh signaling in the developing mammalian thalamus. *Development.* 2012 Aug;139(15):2692-702. PMID: [22745311](https://pubmed.ncbi.nlm.nih.gov/22745311/) **IHC-F (mouse)**

Related products

74-116 Anti-Dlx2 antibody, guinea pig serum