

Anti-HHV6 gQ1 antibody, mouse monoclonal (119)

Product code	65-200
Size	50 µg
Storage	-20°C
Concentration	1.0 mg/ml
Buffer	PBS ⁻ with 50% glycerol
Purity	Purified IgG fraction with protein A from hybridoma cell culture medium.
Immunogen	His6-tagged recombinant gQ1 of HHV-6A encoding 3-422 amino acids expressed in <i>E. coli</i> .
Isotype	Mouse IgG1κ
Reactivity	gQ1 of HHV6A and HHV6B. However, this antibody is not recommended for IP of HHV-6B due to conformation specificity.
Special notes	N/A
Application	<ol style="list-style-type: none"> 1. Western blotting (1/500~1/1,000 dilution) 2. Immunoprecipitation (assay dependent) 3. Immunofluorescence staining and Immunocytochemistry (1/100~1/3,200 dilution) 4. Flow Cytometry (1/100) 5. ELISA (assay dependent)
Background	<p>Human herpesvirus 6 (HHV-6) is the common collective name for Human herpesvirus 6A (HHV-6A) and Human herpesvirus 6B (HHV-6B). These closely related viruses are two of the nine herpesviruses known to have humans as their primary host. HHV-6A and HHV-6B are double stranded DNA viruses within the betaherpesvirinae subfamily and of the genus Roseolovirus. HHV-6A and HHV-6B infects almost all of the human populations tested. The overall nucleotide sequence identity between HHV-6A and HHV-6B is 90% and they are now classified as distinct species.</p> <p>HHV-6A has been described as more neurovirulent, and as such is more frequently found in patients with neuroinflammatory diseases such as multiple sclerosis.</p> <p>HHV-6B primary infection is the cause of the common childhood illness exanthem subitum (also known as roseola infantum or sixth disease). Additionally, HHV-6B reactivation is common in transplant recipients, which can cause several clinical manifestations such as encephalitis, bone marrow suppression and pneumonitis.</p> <p>gQ1 encoded by the U100 gene of HHV6 is glycoprotein, complexes with gH, gL and gQ2 to form HHV6A ligand to CD46 receptor and HHV6B ligand to CD134 receptor. It is expressed in two different forms: an 80-kDa form (gQ1-80K) and a 74-kDa form (gQ1-74K) - only gQ1-80K, but not gQ1-74K, forms the ligand complex with gQ2, gH, and gL. It associates with lipid rafts. Molecular Mass calculated from the gene is 59.9 kDa, 524 amino acids.</p>
Data Link	UniProtKB Q69572 (GQ1_HHV6U)
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

Data Images: 65-200 Anti-HHV6 gQ1 antibody, mouse monoclonal (119)





Fig.1. Identification of gQ1 in HHV-6A infected cells by western blotting using anti-gQ antibody (119). T-cell line HSB-2 cells were infected with HHV-6A at m.o.i of 0.1 and the cells were harvested at 72 h postinfection for lysate preparation for WB. gQ1 is detected as two glycoproteins with 80 kDa and 74 kDa molecular masses.. Sample lysates: (-) Non-treated. (H) Treated with end-glycosylase H. (F) Treated with peptide N-glycosidase F.



1. HSB-2
2. GS-infected HSB-2

Fig.2. Immunoprecipitation of gQ1 from HHV-6A infected cell lysate with anti-gQ1 antibody (119). gQ1 protein was precipitated from the lysate of infected HSB-2 cells by using agarose beads-conjugated anti-gQ1 antibody (119), and processed for western blotting by using anti-gQ1 antibody (119). Lane 1; HSB-2 cells. Lane 2; HSB-2 cells infected with HHV-6A GS strain




Fig.3. Immuno-staining of gQ1 in HHV6A infected HSB-2 cells by using anti-HHV6 gQ1 antibody. The infected cells were harvested 3 days postinfection, fixed in cold acetone and immunostained with FITC-conjugated anti HHV6 gQ1 antibody (119). Specific immunofluorescence was observed with a confocal laser scanning.




Fig.4. Flow cytometry analysis of gQ1 expression on the cell surface in a transient expression system. 293T cells were transfected with gQ1 expression plasmid. The cells were harvested on second day postinfection, fixed with 4% PFA, permeabilized with 0.1% Triton-X100, incubated with anti-gQ1 antibody (119) and then with FITC-conjugated anti-mouse IgG antibody. Histograms show fluorescence intensities measured in arbitrary units on a log scale (*x* axis) and relative cell numbers on a linear scale (*y* axis). Black line is control and red line is gQ1 introduced cells.

References: This antibody has been described in Ref. 1 and used in Ref.1-7.

1. Mori Y. et al. (2003) The Human Herpesvirus 6 U100 Gene Product Is the Third Component of the gH-gL Glycoprotein Complex on the Viral Envelope. *J. Virol.* 77: 2452-2458. [PubMed 12551983](#) **WB, IP**
2. Mori Y. et al (2003) Human Herpesvirus 6 Variant A Glycoprotein H-Glycoprotein L-Glycoprotein Q Complex Associates with Human CD46. *J. Virol.* 77: 4992-4999. [PubMed 12663806](#) **WB, IP**
3. Mori Y. et al. (2004) Discovery of a second form of tripartite complex containing gH-gL of human herpesvirus 6 and observations on CD46. *J Virol.* 78:4609-16. [PubMed 15078943](#). **WB, IP**
4. Akkapaiboon P. et al. (2004) Intracellular processing of human herpesvirus 6 glycoproteins Q1 and

Q2 into tetrameric complexes expressed on the viral envelope. J Virol. 78:7969-83. [PubMed 15254169](#)

WB, IP, IF

5. Huang H. et al (2006) Human herpesvirus 6 envelope cholesterol is required for virus entry. J Gen Virol. 87: 277-285. [PubMed 16432012](#) **WB, Flow Cyt.**

6. Tang H. et al. (2010) Human herpesvirus 6 encoded glycoprotein Q1 gene is essential for virus growth. Virology. 407:360-7. [PubMed 20863544](#) **WB**

7. Tang H. et al (2011) Human herpesvirus 6 glycoprotein complex formation is required for folding and trafficking of the gH/gL/gQ1/gQ2 complex and its cellular receptor binding. J Virol. 85:11121-30. [PubMed 21849437](#) **WB, IP, Flow Cyt.**